



*Committed to children's growth, health,
and development.*

Motrin (ibuprofen) Dosage Chart

Give the weight appropriate dosage listed below every 6 to 8 hours as needed for a temperature > 101 °F or for pain — but should not be given more than four times in twenty-four hours

(Note: Don't use household teaspoons, which can vary in size.)

****DO NOT GIVE IBUPROFEN IF CHILD IS LESS THAN 6 MONTHS OF AGE.****

Weight (lbs)	Infant Drops 50 mg/1.25 ml	Children's 100mg/5ml
8-9	N/A	N/A
10-11	N/A	N/A
12-13	N/A	N/A
14-16	70 mg (1.75 ml)	N/A
17-18	80 mg (2.0 ml)	N/A
19-22	90 mg (2.25 ml)	N/A
22-26	100 mg (2.5 ml)	5ml (1 tsp)
28-31	120 mg (3.0 ml)	6 ml
32-35	140 mg (3.5 ml)	7 ml (1 1/2 tsp)
36-39	160 mg (4.0 ml)	8 ml
40-44	N/A	9 ml
45-48	N/A	10 ml
49-53	N/A	11 ml
54-57	N/A	12 ml
58-62	N/A	13 ml
63-66	N/A	14 ml
67-71	N/A	15 ml
72-75	N/A	16 ml
76-80	N/A	17 ml (3 1/2 tsp)
81-84	N/A	18 ml
85-88	N/A	19 ml
89-93	N/A	20 ml (4 tsp)
94-99	N/A	20 ml (4 tsp)

When used together, Tylenol and motrin may be alternated every 3 hours. Consult a physician before using mortrin if your child has any kidney problems



*Committed to children's growth, health,
and development.*

Tylenol (acetaminophen) Dosage Chart

Give the weight appropriate dosage listed below every 4 hours as needed for a temperature > 101 °F — but should not be given more than five times in twenty-four hours.

(Note: Don't use household teaspoons, which can vary in size.)

****DO NOT GIVE TYLENOL TO CHILDREN UNDER 3 MONTHS OF AGE WITHOUT FIRST CONSULTING A DOCTOR.****

Weight (lbs)	Drops 80 mg per dropper/0.8 ml	Elixir (160 mg per teaspoon/5ml)	Chewable Tablets (80 mg tablets)	Junior Strength (160 mg tablet/capsule)
6-7	40mg (0.4ml)	N/A	N/A	NA/
8-9	40 mg (0.4 ml)	N/A	N/A	N/A
10-11	80 mg (0.8 ml)	N/A	N/A	N/A
12-13	80 mg (0.8 ml)	N/A	N/A	N/A
14-16	80 mg (0.8 ml)	N/A	N/A	N/A
17-18	120 mg (1.2 ml)	N/A	N/A	N/A
19-22	120 mg (1.2 ml)	N/A	N/A	N/A
22-26	160 mg (1.6 ml)	5 ml (1 tsp)	2 tabs	1 tab/cap
28-31	N/A	5 ml	2 tabs	1 tab/cap
32-35	N/A	7 ml (1 1/2 tsp)	3 tabs	1 1/2 tabs/caps
36-39	N/A	7 ml	3 tabs	1 1/2 tabs/caps
40-44	N/A	7 ml	3 tabs	1 1/2 tabs/caps
45-48	N/A	10 ml (2 tsp)	4 tabs	2 tabs/caps
49-53	N/A	10 ml	4 tabs	2 tabs/caps
54-57	N/A	12 ml (2 1/2 tsp)	5 tabs	2 1/2 tabs/caps
59-62	N/A	12 ml	5 tabs	2 1/2 tabs/caps
63-66	N/A	14 ml	5 tabs	2 1/2 tabs/caps
67-71	N/A	15 ml (3 tsp)	6 tabs	3 tabs/caps
72-75	N/A	15 ml	6 tabs	3 tabs/caps
76-80	N/A	17 ml (3 1/2 tsp)	7 tabs	3 1/2 tabs/caps
81-84	N/A	17 ml	7 tabs	3 1/2 tabs/caps
85-88	N/A	17 ml	7 tabs	3 1/2 tabs/caps
89-93	N/A	20 ml (4 tsp)	8 tabs	4 tabs/caps

When used together, Tylenol and motrin may be alternated every 3 hours. Consult a physician before using Tylenol if your child has any liver problems.